VERSAR



DISINFECTANT FOR THE FOOD INDUSTRY

- Bactericidal, Fungicidal, Yeasticidal, Virucidal
- Broad spectrum surface disinfectant for use in the food industry
- Sodium hypochlorite based
- Active against a wide range of pathogens including Candida Albicans and Clostridium Difficile

USES

A highly effective disinfectant concentrate which can be used on a range of hard surfaces, including general hard surfaces such as floors and walls.

May be used as a disinfectant in the milk and meat industry against a broad range of organisms.

Can also be used to soak equipment (non-metal) such as chopping boards, containers, plastic utensils, etc.

DIRECTIONS

For general disinfection, dilute 1 part Versan with 5 parts water. For efficacy against viruses/bacteria, dilute 1 part Versan with 24 parts water. For efficacy against bactericidal spores, use neat.

Application rate: 50 g/m³ once or twice daily in general use.

1-2 times/hour in busy areas.

Fields of use: Indoor and/or outdoor.

For surfaces which are heavily soiled, first clean with a degreaser, such as UBIK 2000 and rinse thoroughly. Apply 'Versan' at the correct dilution.

Allow 5 minute contact time (30 minutes for sporicidal action). Wash away with potable water.





Use disinfectants safely. Always read the label and product information before use. Store in a dry frost-free place at 5-30°C and not in direct sunlight. Store in original container. Dispose of as normal industrial waste. The shelf life of this product is 6 months.

DILUTIONS


Contains: Sodium hypochlorite 2.7% w/w


May fade coloured surfaces. Test on an inconspicuous area first.



Code: 260







5L




DISINFECTANT FOR THE FOOD INDUSTRY


BACTERICIDAL EFFICACY		
TEST	ORGANISMS	TEST CONDITIONS
95 EN 1276	Enterococcus hirae, Escherichia coli, Pseudomonas aeruginosa, Staphylococcus aureus.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes
13697 13697	Staphylococcus aureus, Enterococcus hirae, Escherichia coli, Pseudomonas aeruginosa.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes

YEASTICIDAL/FUNGICIDAL EFFICACY			
TEST	ORGANISMS	TEST CONDITIONS	
1650 ASSES	Aspergillus brasiliensis, Candida albicans.	Clean, 20°C, 5 Minutes Dirty, 20°C, 30 Minutes	
13697 13697	Candida albicans, Aspergillus brasiliensis.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes	

SPORICIDAL EFFICACY PHASE			
TEST	ORGANISMS	TEST CONDITIONS	
13704 13704	Bacillus cereus, Bacillus subtilis, Clostridium difficile.	Dirty (3,0 g/L bovine albumin + 3,0 ml/L sheep erythrocytes), 20 °C, 30 min	









DISINFECTANT FOR HEALTHCARE

- Bactericidal, Fungicidal, Yeasticidal, Virucidal
- Broad spectrum surface disinfectant for use in the healthcare industry
- Sodium hypochlorite based
- Active against a wide range of pathogens including Staphylococcus Aureus and Poliovirus Type 1

USES

A highly effective disinfectant concentrate which can be used on a range of hard surfaces, including general hard surfaces such as floors and walls, worktops, counters, etc. in hospitals, care homes and other healthcare establishments.

Also effective for instrument disinfection (non metal).

DIRECTIONS

For general disinfection, dilute 1 part Versan with 5 parts water. For efficacy against viruses/bacteria, dilute 1 part Versan with 24 parts water. For efficacy against bactericidal spores, use neat.

Application rate: 50 g/m³ once or twice daily in general use.

1-2 times/hour in busy areas.

Fields of use: Indoor and/or outdoor.

For surfaces which are heavily soiled, first clean with a degreaser, such as UBIK 2000 and rinse thoroughly. Apply 'Versan' at the correct dilution.

Allow 5 minute contact time (15 minutes for action against Parvovirus). Wash away with potable water.





Use disinfectants safely. Always read the label and product information before use. Store in a dry frost-free place at 5-30°C and not in direct sunlight. Store in original container. Dispose of as normal industrial waste. The shelf life of this product is 6 months.

DILUTIONS


Contains: Sodium hypochlorite 2.7% w/w

May fade coloured surfaces. Test on an inconspicuous area first.











DISINFECTANT FOR HEALTHCARE

BACTERICIDAL EFFICACY		
TEST	ORGANISMS	TEST CONDITIONS
1276 2489EQ	Enterococcus hirae, Escherichia coli, Pseudomonas aeruginosa, Staphylococcus aureus.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes
13697 13697	Staphylococcus aureus, Enterococcus hirae, Escherichia coli, Pseudomonas aeruginosa.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes
95 EV 13727 9459E0	Enterococcus hirae, Escherichia coli, Pseudomonas aeruginosa, Staphylococcus aureus.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes
05 EN 14561 ASSEQ	Staphylococcus aureus, Enterococcus hirae, Pseudomonas aeruginosa.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes


YEASTICIDAL/FUNGICIDAL EFFICACY			
TEST	ORGANISMS	TEST CONDITIONS	
1650 2495E	Aspergillus brasiliensis, Candida albicans.	Clean, 20°C, 5 Minutes Dirty, 20°C, 30 Minutes	
13624 13624	Aspergillus brasiliensis, Candida albicans.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes	
13697 13697	Candida albicans, Aspergillus brasiliensis.	Clean, 20°C, 5 Minutes Dirty, 20°C, 5 Minutes	
14562 14562	Candida albicans, Aspergillus brasiliensis.	Clean, 20°C, 5 Minutes	

VIRUCIDAL EFFICACY			
TEST	ORGANISMS	TEST CONDITIONS	
14476 14476	Adenovirus Type 5, Murine norovirus, Poliovirus Type 1.	Clean, 20°C, 5 Minutes Dirty, 20°C, 30 Minutes	
05 EN 14476 1455E	Parvovirus.	Clean, 40°C, 15 Minutes Dirty, 40°C, 15 Minutes	

SPORICIDAL EFFICACY PHASE		
TEST	ORGANISMS	TEST CONDITIONS
13704 13704	Bacillus cereus, Bacillus subtilis, Clostridium difficile.	Dirty (3,0 g/L bovine albumin + 3,0 ml/L sheep erythrocytes), 20 °C, 30 min









ERSA



DISINFECTANT FOR VETERINARY HYGIENE

- Bactericidal, Fungicidal, Yeasticidal, Virucidal
- Broad spectrum surface disinfectant for use in the animal care industry
- Sodium hypochlorite based
- Active against a wide range of pathogens including Candida Albicans and Bovine Enterovirus Type 1

USES

A highly effective disinfectant concentrate which can be used on a range of hard surfaces, including general hard surfaces such as floors and walls.

Also effective for the disinfection of surfaces within vehicles which have been used to transport animals.

DIRECTIONS

For general disinfection, dilute 1 part Versan with 5 parts water. For efficacy against viruses/bacteria, dilute 1 part Versan with 24 parts water.

Application rate: 50 g/m³ once or twice daily in general use.

1-2 times/hour in busy areas.

Fields of use: Indoor and/or outdoor.

For surfaces which are heavily soiled, first clean with a degreaser, such as UBIK 2000 and rinse thoroughly. Apply 'Versan' at the correct dilution.

Allow 5 minute contact time (30 minutes for sporicidal action). Wash away with potable water.





Use disinfectants safely. Always read the label and product **information before use.** Store in a dry frost-free place at 5-30°C and not in direct sunlight. Store in original container. Dispose of as normal industrial waste. The shelf life of this product is 6 months.

DILUTIONS

Contains: Sodium hypochlorite 2.7% w/w


May fade coloured surfaces. Test on an inconspicuous area first.



Code: 260







5L



DISINFECTANT FOR VETERINARY HYGIENE

BACTERICIDAL EFFICACY			
TEST	ORGANISMS	TEST CONDITIONS	
1656 1656	Enterococcus hirae, Proteus hauseri, Pseudomonas aeruginosa, Staphylococcus aureus.	Clean, 10°C, 1 Minutes (3,0 g/L bovine albumin) Dirty, 10°C, 5 Minutes (10 g/L skimmed milk)	
14349 14349	Staphylococcus aureus, Proteus hauseri, Enterococcus hirae, Pseudomonas aeruginosa.	Clean, 10°C, 5 Minutes (3,0 g/L bovine albumin) Dirty, 10°C, 5 Minutes (10 g/L skimmed milk)	

YEASTICIDAL/FUNGICIDAL EFFICACY		
TEST	ORGANISMS	TEST CONDITIONS
05 EA 1657 2455ES	Aspergillus brasiliensis, Candida albicans.	Dirty, 10°C, 5 Minutes (10 g/L skimmed milk)

VIRUCIDAL EFFICACY			
TEST	ORGANISMS	TEST CONDITIONS	
14675 14675	bovine enterovirus type 1 - ECBO virus strain LCR-4 (ATCC VR-248)	Dirty, 10°C, 5 Minutes (10 g/L skimmed milk)	





